

Course: Calculus (3)

Chapter: [14]

MULTIPLE INTEGRALS

Section: [14.1]

DOUBLE INTEGRALS




# THE AREA PROBLEM

Given a function  $f$  that is continuous and nonnegative on an interval  $[a, b]$ , find the area between the graph of  $f$  and the interval  $[a, b]$  on the  $x$ -axis.

$$A \approx \sum_{k=1}^n f(x_k^*) \Delta x_k$$

$$A = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*) \Delta x_k$$



$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*) \Delta x_k$$



## THE VOLUME PROBLEM

Given a function  $f$  of two variables that is continuous and nonnegative on a region  $R$  in the  $xy$ –plane, find the volume of the **solid enclosed between the surface  $z = f(x, y)$  and the region  $R$ .**

- Using lines parallel to the coordinate axes, divide the rectangle enclosing the region  $R$  into sub-rectangles, and exclude from consideration all those sub-rectangles that contain any points outside of  $R$ .




# THE VOLUME PROBLEM

$$V \approx \sum_{k=1}^n f(x_k^*, y_k^*) \Delta A_k$$

$$V = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*, y_k^*) \Delta A_k$$

$$\iint_R f(x, y) dA = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*, y_k^*) \Delta A_k$$



which is called the double integral of  $f(x, y)$  over  $R$ .

## EVALUATING DOUBLE INTEGRALS

- The partial derivatives of a function  $f(x, y)$  are calculated by holding one of the variables fixed and differentiating with respect to the other variable.
- Let us consider the reverse of this process, **partial integration**.

$$\int_a^b f(x, y) dx$$

✓ **The partial definite integral with respect to  $x$ .**

✓ Is **evaluated by** holding  $y$  fixed and integrating with respect to  $x$ .

$$\int_c^d f(x, y) dy$$

✓ **The partial definite integral with respect to  $y$ .**

✓ Is **evaluated by** holding  $x$  fixed and integrating with respect to  $y$ .

## EVALUATING DOUBLE INTEGRALS

**Example** (1)  $\int_0^1 xy^2 dx = y^2 \int_0^1 x dx = \frac{y^2 x^2}{2} \Big|_0^1 = \frac{y^2}{2}$

(2)  $\int_0^1 xy^2 dy = x \int_0^1 y^2 dy = \frac{xy^3}{3} \Big|_0^1 = \frac{x}{3}$

- NOTE**
- A partial definite integral with respect to  $x$  is a function of  $y$  and hence can be integrated with respect to  $y$ .
  - A partial definite integral with respect to  $y$  can be integrated with respect to  $x$ .
  - This two-stage integration process is called **iterated** (or *repeated*) **integration**.

# EVALUATING DOUBLE INTEGRALS

- We introduce the following notation:

$$\int_c^d \int_a^b f(x, y) dx dy = \int_c^d \left[ \int_a^b f(x, y) dx \right] dy$$

$$\int_a^b \int_c^d f(x, y) dy dx = \int_a^b \left[ \int_c^d f(x, y) dy \right] dx$$

- These integrals are called *iterated integrals*.

## EVALUATING DOUBLE INTEGRALS

**Example** Evaluate  $\int_1^3 \int_2^4 (40 - 2xy) dy dx$

$$\begin{aligned} \int_1^3 \int_2^4 (40 - 2xy) dy dx &= \int_1^3 \left[ \int_2^4 (40 - 2xy) dy \right] dx \\ &= \int_1^3 (40y - xy^2) \Big|_2^4 dx \\ &= \int_1^3 [(160 - 16x) - (80 - 4x)] dx = \int_1^3 (80 - 12x) dx \\ &= 112 \end{aligned}$$



# EVALUATING DOUBLE INTEGRALS

**Homework** Evaluate  $\int_2^4 \int_1^3 (40 - 2xy) dx dy = 112$


## Fubini's Theorem

Let  $R$  be the rectangle defined by

$$R = \{(x, y) : a \leq x \leq b, c \leq y \leq d\}$$
$$= [a, b] \times [c, d]$$

If  $f(x, y)$  is continuous on this rectangle, then

$$\iint_R f(x, y) dA = \int_a^b \int_c^d f(x, y) dy dx = \int_c^d \int_a^b f(x, y) dx dy$$



## EVALUATING DOUBLE INTEGRALS

**Example** Use a double integral to find the volume of the solid that is bounded above by the plane  $z = 4 - x - y$  and below by the rectangle  $R = [0, 1] \times [0, 2]$ .


$$\begin{aligned} V &= \iint_R (4 - x - y) dA = \int_0^1 \int_0^2 (4 - x - y) dy dx = \int_0^1 \left[ \int_0^2 (4 - x - y) dy \right] dx \\ &= \int_0^1 \left( 4y - xy - \frac{y^2}{2} \right) \Big|_0^2 dx \\ &= \int_0^1 (6 - 2x) dx = 5 = \int_0^2 \int_0^1 (4 - x - y) dx dy \end{aligned}$$

## PROPERTIES OF DOUBLE INTEGRALS

$$\iint_R cf(x, y)dA = c \iint_R f(x, y)dA \quad (c \text{ constant})$$

$$\iint_R [f(x, y) \pm g(x, y)]dA = \iint_R f(x, y)dA \pm \iint_R g(x, y)dA$$

$$\iint_R f(x, y)dA = \iint_{R_1} f(x, y)dA + \iint_{R_2} f(x, y)dA$$



## PROPERTIES OF DOUBLE INTEGRALS

**NOTE** If  $R = [a, b] \times [c, d]$  is a rectangular region, and  $f(x, y) = g(x)h(y)$ , then

$$\iint_R f(x, y) dA = \iint_R g(x)h(y) dA = \left[ \int_a^b g(x) dx \right] \left[ \int_c^d h(y) dy \right]$$

**Example** 
$$\int_0^1 \int_0^2 e^{x+y} dx dy = \int_0^1 \int_0^2 e^x e^y dx dy$$
$$= \left( \int_0^2 e^x dx \right) \left( \int_0^1 e^y dy \right) = (e^2 - 1)(e - 1)$$

## EXERCISE SET 14.1 QUESTION 33

**Homework** Evaluate the integral by choosing a convenient order of integration:

$$\frac{1}{3\pi} = \iint_R x \cos(xy) \cos^2(\pi x) dA \quad ; \quad R = \left[0, \frac{1}{2}\right] \times [0, \pi]$$

Course: Calculus (3)

Chapter: [14]

MULTIPLE INTEGRALS

Section: [14.2]

DOUBLE INTEGRALS OVER NONRECTANGULAR REGIONS

## ITERATED INTEGRALS WITH NONCONSTANT LIMITS OF INTEGRATION

In this section we will see that double integrals over nonrectangular regions can often be evaluated as iterated integrals

**Example** 
$$\int_0^1 \int_{-x}^{x^2} y^2 x \, dy dx = \int_0^1 \left[ \int_{-x}^{x^2} y^2 x \, dy \right] dx = \int_0^1 \left. \frac{xy^3}{3} \right|_{-x}^{x^2} dx$$
$$= \int_0^1 \left( \frac{x^7}{3} + \frac{x^4}{3} \right) dx = \left( \frac{x^8}{24} + \frac{x^5}{15} \right) \Big|_0^1 = \frac{13}{120}$$

# ITERATED INTEGRALS WITH NONCONSTANT LIMITS OF INTEGRATION

**Example**  $\int_0^{\pi/3} \int_0^{\cos y} x \sin y \, dx dy = \int_0^{\pi/3} \left[ \int_0^{\cos y} x \sin y \, dx \right] dy$

By Substitution

Let  $t = \cos y$

$$\frac{dt}{dy} = -\sin y$$

$$dy = -\frac{dt}{\sin y}$$

$$y = \pi/3$$

$$t = 1/2$$

$$y = 0$$

$$t = 1$$


$$\begin{aligned} &= \int_0^{\pi/3} \left. \frac{x^2 \sin y}{2} \right|_0^{\cos y} dy \\ &= \int_0^{\pi/3} \frac{1}{2} \cos^2 y \sin y \, dy = -\frac{1}{2} \int_1^{1/2} t^2 \sin y \frac{dt}{\sin y} \\ &= \frac{1}{2} \int_{1/2}^1 t^2 dt = \left. \frac{t^3}{6} \right|_{1/2}^1 = \frac{7}{48} \end{aligned}$$



# DOUBLE INTEGRALS OVER NONRECTANGULAR REGIONS


## Type I Region

is bounded on the left and right by vertical lines  $x = a$  and  $x = b$  and is bounded below and above by continuous curves  $y = g_1(x)$  and  $y = g_2(x)$ , where  $g_1(x) \leq g_2(x)$  for  $a \leq x \leq b$ .



## Type II Region

is bounded below and above by horizontal lines  $y = c$  and  $y = d$  and is bounded on the left and right by continuous curves  $x = h_1(y)$  and  $x = h_2(y)$  satisfying  $h_1(y) \leq h_2(y)$  for  $c \leq y \leq d$ .



# DOUBLE INTEGRALS OVER NONRECTANGULAR REGIONS

1) If  $R$  is a **type I region** on which  $f(x, y)$  is continuous, then

$$\iint_R f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx$$

2) If  $R$  is a **type II region** on which  $f(x, y)$  is continuous, then


$$\iint_R f(x, y) dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy$$

## DOUBLE INTEGRALS OVER NONRECTANGULAR REGIONS

**Example** Evaluate  $\iint_R xy dA$  over the region  $R$  enclosed between  $y = \frac{1}{2}x$ ,  $y = \sqrt{x}$ ,  $x = 2$  and  $x = 4$ .

### Type I Region


$$\begin{aligned}\iint_R xy dA &= \int \int xy dy dx = \int_2^4 \left[ \int_{x/2}^{\sqrt{x}} xy dy \right] dx \\ &= \int_2^4 \left. \frac{xy^2}{2} \right|_{x/2}^{\sqrt{x}} dx = \int_2^4 \left( \frac{x^2}{2} - \frac{x^3}{8} \right) dx = \frac{11}{6}\end{aligned}$$



## DOUBLE INTEGRALS OVER NONRECTANGULAR REGIONS

**Example** Evaluate  $\iint_R (2x - y^2) dA$  over the triangular region  $R$  enclosed between the lines  $y = -x + 1$ ,  $y = x + 1$ , and  $y = 3$ .


$$\begin{aligned}\iint_R (2x - y^2) dA &= \int \int (2x - y^2) dx dy \\ &= \int_1^3 x^2 - xy^2 \Big|_{1-y}^{y-1} dy \\ &= \int_1^3 (2y^2 - 2y^3) dy = -\frac{68}{3}\end{aligned}$$



## DOUBLE INTEGRALS OVER NONRECTANGULAR REGIONS

**Example** Evaluate  $\iint_R (2x - y^2) dA$  over the triangular region  $R$  enclosed between the lines  $y = -x + 1$ ,  $y = x + 1$ , and  $y = 3$ .

$$\iint_R (2x - y^2) dA$$




## DOUBLE INTEGRALS OVER NONRECTANGULAR REGIONS

**Example** Evaluate  $\iint_R (2x - y^2) dA$  over the triangular region  $R$  enclosed between the lines  $y = -x + 1$ ,  $y = x + 1$ , and  $y = 3$ .

$$\iint_R (2x - y^2) dA = \iint_{R_1} (2x - y^2) dA + \iint_{R_2} (2x - y^2) dA$$

$$= \int_{-2}^0 \int_{-x+1}^3 (2x - y^2) dy dx + \int_0^2 \int_{x+1}^3 (2x - y^2) dy dx$$




## DOUBLE INTEGRALS OVER NONRECTANGULAR REGIONS

**Example** Evaluate  $\int_0^2 \int_{y/2}^1 e^{x^2} dx dy$   
 $y/2 \rightarrow y = 2x$

Since there is no elementary antiderivative of  $e^{x^2}$ , the integral cannot be evaluated by performing the  $x$  –integration first.

We will try to evaluate this integral by expressing it as an equivalent iterated integral with the order of integration reversed.



## DOUBLE INTEGRALS OVER NONRECTANGULAR REGIONS


**Example** Evaluate  $\int_0^2 \int_{y/2}^1 e^{x^2} dx dy$

$$\int_0^2 \int_{y/2}^1 e^{x^2} dx dy = \int \int e^{x^2} dy dx = \int_0^1 \left[ \int_0^{2x} e^{x^2} dy \right] dx$$

$$= \int_0^1 e^{x^2} y \Big|_0^{2x} dx$$

By Substitution  
Let  $t = x^2$

$$\Rightarrow \int_0^1 2xe^{x^2} dx = \int_0^1 e^t dt = e - 1$$






## AREA CALCULATED AS A DOUBLE INTEGRAL

### Example

Use a double integral to find the area of the region  $R$  enclosed between the parabola  $y = \frac{1}{2}x^2$  and the line  $y = 2x$ .

$$\text{area of } R = \iint_R 1 \, dA = \iint_R dA$$




Cylinder with base  $R$  and height 1

## AREA CALCULATED AS A DOUBLE INTEGRAL

$$\text{area of } R = \iint_R 1 \, dA = \iint_R dA$$

**Example** Use a double integral to find the area of the region  $R$  enclosed between the parabola  $y = \frac{1}{2}x^2$  and the line  $y = 2x$ .

$$\begin{aligned} \text{Area of } R &= \iint_R dA && \text{(Type II Region)} \\ &= \int_0^8 \int_{y/2}^{\sqrt{2y}} dx dy = \int_0^8 x \Big|_{y/2}^{\sqrt{2y}} dy \\ &= \int_0^8 \left( \sqrt{2y} - \frac{y}{2} \right) dy = \frac{16}{3} \end{aligned}$$




## AREA CALCULATED AS A DOUBLE INTEGRAL

$$\text{area of } R = \iint_R 1 \, dA = \iint_R dA$$

**Example** Use a double integral to find the area of the region  $R$  enclosed between the parabola  $y = \frac{1}{2}x^2$  and the line  $y = 2x$ .

$$\begin{aligned} \text{Area of } R &= \iint_R dA && \text{(Type I Region)} \\ &= \int_0^4 \int_{x^2/2}^{2x} dy dx = \int_0^4 y \Big|_{x^2/2}^{2x} dx \\ &= \int_0^4 \left( 2x - \frac{x^2}{2} \right) dx = \frac{16}{3} \end{aligned}$$




## EXERCISE SET 14.2

9. Let  $R$  be the region shown in the accompanying figure. Fill in the missing limits of integration.

$$(a) \iint_R f(x, y) dA = \int_{\square}^{\square} \int_{\square}^{\square} f(x, y) dy dx$$

$$(b) \iint_R f(x, y) dA = \int_{\square}^{\square} \int_{\square}^{\square} f(x, y) dx dy$$



Course: Calculus (3)

Chapter: [14]

MULTIPLE INTEGRALS

Section: [14.3]

DOUBLE INTEGRALS IN POLAR COORDINATES


## SIMPLE POLAR REGIONS

- Some double integrals are easier to evaluate if the region of integration is expressed in polar coordinates.
- This is usually true if the region is bounded by any curve whose equation is simpler in polar coordinates than in rectangular coordinates.
- **Example:** Consider the quarter-disk  $x^2 + y^2 = 4$  in the first quadrant shown below.

Rectangular  
Coordinates

$$0 \leq x \leq 2$$

$$0 \leq y \leq \sqrt{4 - x^2}$$




Polar  
Coordinates

$$0 \leq r \leq 2$$

$$0 \leq \theta \leq \pi/2$$


## DOUBLE INTEGRALS IN POLAR COORDINATES

**NOTE** A **polar rectangle** is a simple polar region for which the bounding polar curves are circular arcs.



**Theorem** If  $R$  is a simple polar region whose boundaries are the rays  $\theta = \alpha$  and  $\theta = \beta$  and the curves  $r = r_1(\theta)$  and  $r = r_2(\theta)$ , and if  $f(r, \theta)$  is continuous on  $R$ , then


$$\iint_R f(r, \theta) dA = \int_{\alpha}^{\beta} \int_{r_1(\theta)}^{r_2(\theta)} f(r, \theta) r dr d\theta$$



## DOUBLE INTEGRALS IN POLAR COORDINATES

**Example** Find the volume of the solid bounded by the cylinder  $x^2 + y^2 = 4$  and the plane  $y + z = 4$ .


$$\begin{aligned} V &= \iint_R (4 - y) dA = \int \int (4 - r \sin \theta) r dr d\theta \\ &= \int_0^{2\pi} \left[ \int_0^2 (4r - r^2 \sin \theta) dr \right] d\theta \\ &= \int_0^{2\pi} \left( 2r^2 - \frac{1}{3} r^3 \sin \theta \right) \Big|_0^2 d\theta = \int_0^{2\pi} \left( 8 - \frac{8}{3} \sin \theta \right) d\theta \\ &= \left( 8\theta + \frac{8}{3} \cos \theta \right) \Big|_0^{2\pi} = 16\pi \end{aligned}$$





## DOUBLE INTEGRALS IN POLAR COORDINATES

**Example** Evaluate  $\int_{-1}^1 \int_0^{\sqrt{1-x^2}} (x^2 + y^2)^{3/2} dy dx$




$$\begin{aligned} \int_{-1}^1 \int_0^{\sqrt{1-x^2}} (x^2 + y^2)^{3/2} dy dx &= \iint (r^2)^{3/2} r dr d\theta \\ &= \int_0^{\pi} \int_0^1 r^4 dr d\theta = \int_0^{\pi} \frac{1}{5} d\theta = \frac{\pi}{5} \end{aligned}$$

## DOUBLE INTEGRALS IN POLAR COORDINATES

**Example** Evaluate  $\iint_R \frac{1}{1+x^2+y^2} dA$  where  $R$  is the region in the first quadrant bounded by  $y = 0$ ,  $y = x$ ,  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 4$ .

$$\iint_R \frac{1}{1+x^2+y^2} dA = \int \int \frac{1}{1+r^2} r dr d\theta$$

$$= \int_0^{\pi/4} \left[ \int_1^2 \frac{r}{1+r^2} dr \right] d\theta$$




$$\tan \theta = \frac{y}{x} = \frac{x}{x} = 1$$

$$\theta = \frac{\pi}{4}$$

## DOUBLE INTEGRALS IN POLAR COORDINATES

**Example** Evaluate  $\iint_R \frac{1}{1+x^2+y^2} dA$  where  $R$  is the region in the first quadrant bounded by  $y = 0$ ,  $y = x$ ,  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 4$ .




$$\tan \theta = \frac{y}{x} = \frac{x}{x} = 1$$
$$\theta = \frac{\pi}{4}$$

$$\begin{aligned} \iint_R \frac{1}{1+x^2+y^2} dA &= \int \int \frac{1}{1+r^2} r dr d\theta \\ &= \int_0^{\pi/4} \left[ \frac{1}{2} \int_1^2 \frac{2r}{1+r^2} dr \right] d\theta = \int_0^{\pi/4} \frac{1}{2} \ln|1+r^2| \Big|_1^2 d\theta \\ &= \int_0^{\pi/4} \frac{1}{2} \ln \left( \frac{5}{2} \right) d\theta = \frac{\pi}{8} \ln \left( \frac{5}{2} \right) \end{aligned}$$

## DOUBLE INTEGRALS IN POLAR COORDINATES

**Example** Use a double-integral to show that the area of the region  $R$  shown is  $\frac{9\pi}{2}$ .


$$\text{Area of } R = \iint_R dA = \int \int r dr d\theta$$



## DOUBLE INTEGRALS IN POLAR COORDINATES

**Example** Use a double-integral to show that the area of the region  $R$  shown is  $\frac{9\pi}{2}$ .

$$\begin{aligned}\text{Area of } R &= \iint_R dA = \int \int_0^3 r dr d\theta \\ &= \int_{-\pi/3}^{2\pi/3} \left[ \int_0^3 r dr \right] d\theta = \int_{-\pi/3}^{2\pi/3} \left. \frac{r^2}{2} \right|_0^3 d\theta \\ &= \int_{-\pi/3}^{2\pi/3} \frac{9}{2} d\theta = \frac{9}{2} \theta \Big|_{-\pi/3}^{2\pi/3} = \frac{9\pi}{2}\end{aligned}$$



## DOUBLE INTEGRALS IN POLAR COORDINATES

**Example** Evaluate  $\int_0^{\infty} e^{-x^2} dx = I$

$$\begin{aligned} I^2 &= \left( \int_0^{\infty} e^{-x^2} dx \right)^2 = \left( \int_0^{\infty} e^{-x^2} dx \right) \left( \int_0^{\infty} e^{-x^2} dx \right) \\ &= \left( \int_0^{\infty} e^{-x^2} dx \right) \left( \int_0^{\infty} e^{-y^2} dy \right) \\ &= \int_0^{\infty} \int_0^{\infty} e^{-x^2} e^{-y^2} dx dy = \int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy \end{aligned}$$

## DOUBLE INTEGRALS IN POLAR COORDINATES


**Example** Evaluate  $\int_0^{\infty} e^{-x^2} dx = I$

$$I^2 = \int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy = \int \int e^{-r^2} r dr d\theta$$

$$= \int_0^{\pi/2} \left[ \int_0^{\infty} r e^{-r^2} dr \right] d\theta$$

By substitution. Let  $t = r^2$ .

$$= \int_0^{\pi/2} \left[ \int_0^{\infty} \frac{1}{2} e^{-t} dt \right] d\theta = \int_0^{\pi/2} \left. \frac{-1}{2} e^{-t} \right|_0^{\infty} d\theta = \int_0^{\pi/2} \frac{1}{2} d\theta = \frac{\pi}{4}$$



Course: Calculus (3)

Chapter: [14]

MULTIPLE INTEGRALS

Section: [14.5]

Triple Integral [Iterated Method]



## EVALUATING TRIPLE INTEGRALS OVER RECTANGULAR BOXES

Let  $G$  be the rectangular box defined by the inequalities

$$a \leq x \leq b \quad , \quad c \leq y \leq d \quad , \quad k \leq z \leq \ell$$

If  $f$  is continuous on the region  $G$ , then

$$\iiint_G f(x, y, z) dV = \int_a^b \int_c^d \int_k^\ell f(x, y, z) dz dy dx$$

Moreover, the iterated integral on the right can be replaced with any of the five other iterated integrals that result by altering the order of integration.

## EVALUATING TRIPLE INTEGRALS OVER RECTANGULAR BOXES

**Example** Evaluate the triple integral  $\iiint_G 12xy^2z^3 dV$  over the rectangular box  
 $G = [-1,2] \times [0,3] \times [0,2]$

$$\iiint_G 12xy^2z^3 dV = \int_{-1}^2 \int_0^3 \int_0^2 12xy^2z^3 dz dy dx = \int_{-1}^2 \int_0^3 \left[ \int_0^2 12xy^2z^3 dz \right] dy dx$$

$$= \int_{-1}^2 \int_0^3 48xy^2 dy dx = \int_{-1}^2 432x dx = 648$$

$$\iiint_G 12xy^2z^3 dV = 12 \left[ \int_{-1}^2 x dx \right] \left[ \int_0^3 y^2 dy \right] \left[ \int_0^2 z^3 dz \right] = 648$$

## EVALUATING TRIPLE INTEGRALS OVER MORE GENERAL REGIONS

**Example** Evaluate  $\int_0^1 \int_0^y \int_0^{\sqrt{1-y^2}} z \, dz dx dy$

$$\begin{aligned} \int_0^1 \int_0^y \int_0^{\sqrt{1-y^2}} z \, dz dx dy &= \int_0^1 \int_0^y \left. \frac{1}{2} z^2 \right|_0^{\sqrt{1-y^2}} dx dy = \int_0^1 \int_0^y \frac{1}{2} (1 - y^2) dx dy \\ &= \int_0^1 \left. \frac{1}{2} (1 - y^2) x \right|_0^y dy = \int_0^1 \frac{1}{2} (1 - y^2) y dy \\ &= \frac{1}{2} \int_0^1 (y - y^3) dy = \frac{1}{8} \end{aligned}$$